

**Supplementary Data 1.** Additional Analysis of Percentage of Correct Answers for Functional Health Literacy Test and Nutrition Labeling Health Literacy Test

Items	Total (n=46)		LHL (n=30)		HHL (n=16)	
	Correct answer rate (%)	Rank	Correct answer rate (%)	Rank	Correct answer rate (%)	Rank
<b>Functional health literacy test</b>						
<b>A. Taking medication</b>						
1. Medicine to prevent blood clots	54.4	1	46.7	2	62.5	1
2. When to take nitroglycerin	19.6	6	6.7	4	37.5	9
3. Dose after missed last dose	50.0	2	53.3	1	43.8	8
<b>B. Laboratory results</b>						
4. Classification of tested body mass index	15.2	10	6.7	4	31.3	10
5. Normality of tested blood sugar	34.8	3	16.7	3	62.5	1
6. Classification of tested total cholesterol	19.6	6	0.0	10	56.3	3
7. Normality of tested HDL-cholesterol	19.6	6	3.3	7	50.0	5
<b>Nutrition labeling health literacy test</b>						
1. Calculate percentage of calories per day	21.7	4	6.7	4	50.0	5
2. Percentage of sodium eaten compared to the recommended daily intake of sodium	21.7	4	3.3	7	56.3	3
3. Calculate the amount of total saturated fat consumed for the day	19.6	6	3.3	7	50.0	5

HDL=high density lipoprotein; HHL=high health literacy; LHL=low health literacy.